

Worldwide Survey of the “Assessing Pain, Both Spontaneous Awakening and Breathing Trials, Choice of Drugs, Delirium Monitoring/Management, Early Exercise/Mobility, and Family Empowerment” (ABCDEF) Bundle

Alessandro Morandi, MD, MPH^{1,2}; Simone Piva, MD³; E. Wesley Ely, MD, MPH⁴; Sheila Nainan Myatra, MD, FCCM, FICCM⁵; Jorge I.F. Salluh, MD, PhD⁶; Dawit Amare, MD⁷; Elie Azoulay, MD, PhD⁸; Giuseppe Bellelli, MD^{2,9}; Akos Csomos, MD, PhD¹⁰; Eddy Fan, MD, PhD^{11,12}; Nazzareno Fagoni, MD, PhD³; Timothy D. Girard, MD, MSCI¹³; Gabriel Heras La Calle, MD¹⁴; Shigeaki Inoue, MD, PhD, MSCI¹⁵; Chae-Man Lim, MD, PhD, FCCM¹⁶; Rafael Kaps, MD¹⁷; Katarzyna Kotfis, MD, PhD¹⁸; Younsuck Koh, MD¹⁶; David Misango, MD¹⁹; Pratik P. Pandharipande, MD, MSCI²⁰; Chairat Permpikul, MD²¹; Cheng Cheng Tan, MD, PhD²²; Dong-Xin Wang, MD, PhD²²; Tarek Sharshar, MD, PhD²³; Yahya Shehabi, PhD, FCICM, FANZCA, EMBA^{24,25}; Yoanna Skrobik, MD, FRCP(c) MSc, FCCM^{26,27}; Jeffrey M. Singh, MD^{28,29}; Arjen Slooter, MD, PhD³⁰; Martin Smith, MBBS, FRCA, FFICM³¹; Ryosuke Tsuruta, MD, PhD³²; Nicola Latronico, MD^{33,34}

¹Department of Rehabilitation, Ancelle Hospital, Cremona, Italy.

²Geriatric Research Group, Brescia, Italy.

³Department of Anesthesia, Critical Care and Emergency, Spedali Civili University Hospital, Brescia, Italy.

⁴Division of Pulmonary and Critical Care and Health Services Research, Vanderbilt University and VA Geriatric Research Education Clinical Center (GRECC), Nashville, TN.

⁵Department of Anaesthesiology, Critical Care and Pain, Tata Memorial Hospital, Mumbai, India.

⁶Department of Critical Care, D'or Institute for Research and Education and Post-Graduate Program Federal University of Rio de Janeiro, Rio de Janeiro, Brazil.

⁷Addis Hiwot General Hospital, Addis Ababa, Ethiopia.

⁸Medical Intensive Care Unit, Hôpital Saint-Louis, ECSTRA team, Biostatistics and clinical epidemiology, UMR 1153 (Center of Epidemiology and Biostatistics Sorbonne Paris Cité, CRESS), INSERM, Paris Diderot Sorbonne University, Paris, France.

⁹School of Medicine and Surgery, University of Milano-Bicocca, Milano Geriatric Clinic, San Gerardo University Hospital, Monza, Italy.

¹⁰Department of Anaesthesia and Intensive Care, Medical Centre, Hungarian Defence Force, Budapest, Hungary.

¹¹Interdepartmental Division of Critical Care Medicine and Institute of Health Policy, Management and Evaluation, University of Toronto, Toronto, ON, Canada.

Copyright © 2017 by the Society of Critical Care Medicine and Wolters Kluwer Health, Inc. All Rights Reserved.

DOI: 10.1097/CCM.0000000000002640

¹² University Health Network, Toronto, ON, Canada.

¹³ CRISMA Center, Department of Critical Care Medicine, University of Pittsburgh and UPMC Health System, Pittsburgh, PA.

¹⁴ International Research Project Humanizing Intensive Care (Proyecto HU-CI), Intensive Care Unit, Hospital Universitario de Torrejón, Madrid, Spain.

¹⁵Department of Emergency and Critical Care Medicine, Tokai University, School of Medicine, Kanagawa, Japan.

¹⁶ Department of Pulmonary and Critical Care Medicine Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea.

¹⁷ General Hospital Novo mesto, Novo mesto, Slovenia.

¹⁸Department of Anaesthesia, Intensive Care and Acute Poisoning, Pomeranian Medical University, Szczecin, Poland.

¹⁹ Aga Khan University Hospital, Nairobi, Kenya.

²⁰ Department of Anesthesiology, Division of Anesthesiology Critical Care Medicine Vanderbilt University Medical Center, Nashville, TN.

²¹ Siriraj Hospital, Bangkok, Thailand.

²² Department of Anesthesiology and Critical Care Medicine, Peking University First Hospital, Beijing, China.

²³Department of Intensive Care Medicine Raymond Poincaré Hospital, Paris, France.

²⁴ School of Clinical Sciences, Faculty of Medicine, Monash University & Medical Center, Melbourne, VIC, Australia.

²⁵ University New South Wales, Clinical School of Medicine, NSW, Australia.

²⁶ Department of Medicine, McGill University, Montreal, QC, Canada.

²⁷ Regroupement de Soins Critiques Respiratoires, Réseau de Santé Respiratoire FRQS, Montreal, QC, Canada.

²⁸Interdepartmental Division of Critical Care Medicine, University of Toronto, Toronto, ON, Canada.

²⁹Krembil Neuroscience Centre, University Health Network, Toronto, ON, Canada.

³⁰Department of Intensive Care Medicine, Brain Center Rudolf Magnus, Utrecht University, University Medical Center Utrecht, The Netherlands.

³¹Department of Neurocritical Care, University College London Hospitals, UCLH/UCL National Institute for Health Research Biomedical Research Centre, London, United Kingdom.

³²Acute & General Medicine, Yamaguchi Graduate School of Medicine, Yamaguchi, Japan.

³³Department of Surgical Specialties, Radiological Sciences and Public Health, University of Brescia, Brescia, Italy.

³⁴Department of Anesthesia, Critical Care and Emergency, Spedali Civili University Hospital, Brescia, Italy.

All authors contributed in study conception; design and critically revised the article; and interpretation of results. Drs. Morandi and Piva have equally contributed as first authors to the article and drafted the article and in acquisition of data.

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's website (<http://journals.lww.com/ccmjjournal>).

Dr. Ely's institution received funding from Veterans Affairs Geriatric Research Education and Clinical Centers and the National Institutes of Health (NIH) (R01AG035117 and R01HL111111); he received funding from Orion, Abbott, and Pfizer; and he received support for article research from the NIH. Dr. Azoulay's institution received research grant funding from Fisher & Paykel, Pfizer, Gilead, and Alexion, and he received funding for board member lectures from Alexion, Astellas, and Gilead. Dr. Fan is supported by a New Investigator Award from the Canadian Institutes of Health Research. Dr. Pandharipande's institution received funding from Hospira Inc in collaboration with the NIH for a research grant. Dr. Pandharipande (R01HL111111 and R01AG035117) and Dr. Needham (R24HL111895 and R24AG054259) are supported by the NIH. Dr. Permpikul disclosed government work. Dr. Shehabi reported related unrestricted research and educational grants from Pfizer and Orion Pharma and the National Health and Medical Research Council of Australia. Speaker's honorarium and travel expenses reimbursed to employing institution. Dr. Slooter works on the development of an EEG-based delirium monitor, any (future) profits of this technology will be used for future scientific research only, and he is supported by the European Society of Intensive Care Medicine, Technology Foundation STW (14066) and the European Union (H2020-PHC-12-2014-672974). The remaining authors have disclosed that they do not have any potential conflicts of interest.

Address requests for reprints to: Alessandro Morandi MD, MPH, Via Aselli 16, Cremona, 26100. E-mail: morandi.alessandro@gmail.com

Objectives: To assess the knowledge and use of the Assessment, prevention, and management of pain; spontaneous awakening and breathing trials; Choice of analgesia and sedation; Delirium assessment; Early mobility and exercise; and Family engagement and empowerment (ABCDEF) bundle to implement the Pain, Agitation, Delirium guidelines.

Design: Worldwide online survey.

Setting: Intensive care.

Intervention: A cross-sectional online survey using the Delphi method was administered to intensivists worldwide, to assess the knowledge and use of all aspects of the ABCDEF bundle.

Measurement and Main Results: There were 1,521 respondents from 47 countries, 57% had implemented the ABCDEF bundle, with varying degrees of compliance across continents. Most of the respondents (83%) used a scale to evaluate pain. Spontaneous awakening trials and spontaneous breathing trials are performed in 66% and 67% of the responder ICUs, respec-

tively. Sedation scale was used in 89% of ICUs. Delirium monitoring was implemented in 70% of ICUs, but only 42% used a validated delirium tool. Likewise, early mobilization was "prescribed" by most, but 69% had no mobility team and 79% used no formal mobility scale. Only 36% of the respondents assessed ICU-acquired weakness. Family members were actively involved in 67% of ICUs; however, only 33% used dedicated staff to support families and only 35% reported that their unit was open 24 hr/d for family visits.

Conclusions: The current implementation of the ABCDEF bundle varies across individual components and regions. We identified specific targets for quality improvement and adoption of the ABCDEF bundle. Our data reflect a significant but incomplete shift toward patient- and family-centered ICU care in accordance with the Pain, Agitation, Delirium guidelines. (*Crit Care Med* 2017; XX:00–00)

Key Words: ABCDEF; intensive care; pain; sedation; spontaneous awakening and breathing trials

Survivors of critical illness often experience persistent physical, mental, and cognitive impairment (1–5). Inadequately treated pain, excessive sedation, delirium, and reduced mobilization have emerged as risk factors for acute muscle wasting and weakness, persisting physical dysfunction and cognitive decline. To aid adoption of the Society of Critical Care Medicine's (SCCM) Pain, Agitation, Delirium (PAD) guidelines, an evidence-based multicomponent and interprofessional team management strategy, known as "Assess, prevent, and manage pain; Both spontaneous awakening trials (SATs) and spontaneous breathing trials (SBTs); attention to the Choice of analgesia and sedation; Delirium monitoring and management; Early mobility and exercise; and Family engagement and empowerment (ABCDEF) bundle" has been developed and implemented in thousands of ICUs (6–9). Each component of the ABCDEF bundle addresses a target practice in the ICU independently associated with patient safety or patient-centered outcomes (10–26). For example, multiple studies have demonstrated the efficacy of implementing combined SATs and SBTs to shorten duration of mechanical ventilation and ICU length of stay (10–13). To date, the ABCDEF bundle approach has been evaluated in only a few countries, and some reports are available on the individual components (27–33). Results vary widely across different countries and jurisdictions (34).

Evaluation of the current state of understanding and implementation of the ABCDEF bundle would help future knowledge translation efforts and identify targets for quality improvement initiatives. We thus employed an international, web-based survey to assess (1) knowledge of the ABCDEF bundle and (2) differences in the use of each component across the world.

METHODS

We conducted a worldwide survey of intensivists (i.e., physicians) endorsed by the European Society of Intensive Care

Medicine (ESICM). The survey instrument described the ABCDEF bundle and then probed eight domains with a total of 41 questions (**supplemental survey**, Supplemental Digital Content 1, <http://links.lww.com/CCM/C792>). Sixty-eight questions were initially created by A.M. and S.P. and were then reduced to 41 using a Delphi method by a steering committee (SC), who were asked to rate each question on a Likert scale ranging from “retain” to “exclude.” The survey was then pretested by the SC, who provided written feedback on “Face validity, Content Validity, and Criterion Validity.” Last, the instrument was “pilot tested” by the SC, evaluating duration, flow, relevance, and acceptability, and questions were screened for redundancy, relevance, and clarity. “Clinical sensitivity testing” was completed by the SC using a seven-question instrument (**Appendix 1**, Supplemental Digital Content 2, <http://links.lww.com/CCM/C793>), and “test-retest reliability” was assessed by the SC who repeated duplicate surveys within 4 weeks.

The local ethics committee reviewed the study and waived the need for approval.

Survey Administration

The survey and a cover letter were distributed to members of ESICM and other national and regional intensive care societies between March 1 and September 15, 2016 via Lime Survey (LimeSurvey GmbH, Hamburg, Germany). To expand survey distribution and improve response rate, SC members sent reminder e-mails with a cover letter, and a web-based advertisement was sent to the following societies: ESICM, the Italian Society of Anesthesia, Analgesia, Reanimation and Intensive Care, the Indian SCCM, the Japanese SCCM, and the Australasian and New Zealand Intensive Care Society.

The survey was open and anonymous, and the database was securely maintained at the University of Brescia (www.anestbs.com). Question sequence was randomized to avoid bias. Date, total time, and single-item time of compilation were recorded. The computer internet protocol address of the respondent was recorded to avoid duplicate entries. The Checklist for Reporting Results of Internet E-Surveys checklist was used to report the data (35).

Data Analysis

Only complete questionnaires were included in the final analysis. Aggregated responses are reported as frequencies and percentages. Continuous data are reported as mean (SD). For test-retest analysis, chi-square test for nominal data, Spearman rho for ordinal data, and Pearson *r* for interval data were used. Analyses were performed with Stata 13.0 (STATA Corp, College Station, TX) software.

RESULTS

We received 1,521 completed questionnaires from respondents in Europe (*n* = 607), South America (*n* = 265), Asia (*n* = 441), North America (*n* = 120), Oceania (*n* = 45), and Africa (*n* = 43). The most represented countries were Italy (*n* = 371; 24%), followed by India (*n* = 250; 16%) and Brazil (*n* = 159,

10%) (**Appendix 2**, Fig. 1, Supplemental Digital Content 2, <http://links.lww.com/CCM/C793>; and **Appendix 3**, Supplemental Digital Content 2, <http://links.lww.com/CCM/C793>). The characteristics of the respondents are listed in **Table 1**. Just over half of respondents reported implementing the PAD guidelines (56.2%) and the ABCDEF bundle (56.6%). Overall, the ABCDEF bundle implementation was greater in non-academic hospitals, in open/semiopen ICUs, and in ICU with larger annual admissions (**Appendix 4**, Supplemental Digital Content 2, <http://links.lww.com/CCM/C793>).

Assess, Prevent, and Manage Pain

Though responses varied geographically, most respondents (83%) reported using a scale to evaluate pain (**Table 2**; and **Appendix 5**, Fig. 2, Supplemental Digital Content 2, <http://links.lww.com/CCM/C793>). The pain scale reported to be most widely used were the Visual Analogic Scale (VAS) (54%) and the Numerical Rating Scale (NRS) (54%). Only 56% of the respondents reported using preemptive analgesia before nursing procedures. The preferred analgesics were morphine (78%), fentanyl (79%), and paracetamol (69%) alone or in combination.

Both SAT and SBT

Two thirds of respondents reported performed SATs, most often once daily (59%) but with wide geographic variation (**Table 2**; and **Appendix 5**, Fig. 2, Supplemental Digital Content 2, <http://links.lww.com/CCM/C793>). Similarly, 67% of respondents reported using SBTs. Only 42% of respondents reported using a coordinated protocol synchronizing SAT and SBT (i.e., a “wake up and breathe” protocol), most often performed by physicians (27%) or nurses (15%).

Choice of Analgesia and Sedation

Respondents (61%) most often reported using the Richmond Agitation Sedation Scale (RASS) to evaluate the level of arousal, followed by the Ramsey scale (22%) (**Table 2**; and **Appendix 5**, Fig. 2, Supplemental Digital Content 2, <http://links.lww.com/CCM/C793>). Just over one third of respondents reported using a sedation protocol. The sedation protocols reported typically (90%) focus on using minimal or no sedation with avoidance of benzodiazepines. When treating an agitated patient, most respondents reported that they evaluate pain first and then delirium before considering using sedation.

Delirium

Half of the respondents estimate that 30% or less of patients in their ICU have delirium (**Table 3**; and **Appendix 5**, Fig. 2, Supplemental Digital Content 2, <http://links.lww.com/CCM/C793>). One third of respondents do not routinely monitor delirium, whereas 40% report that they assess patients for delirium once a day, and 30% report doing so more than once a day, with substantial variation across continents. More than half of the respondents (58%) do not use specific tools to monitor delirium, though they acknowledge the need for delirium monitoring. Among those who use a tool, CAM-ICU is preferred (83%) followed by the ICDSC (17%).

TABLE 1. Characteristics of 1,521 Responders in 47 Countries

Variables	n (%)
Age	
25–35	314 (21)
36–45	585 (38)
46–55	368 (24)
56–65	223 (15)
> 65	20 (2)
Years of clinical experience	
5–10	399 (32)
11–20	476 (40)
> 20	365 (29)
Specialty	
Anesthesiology	603 (40)
Critical care	488 (32)
Internal medicine	190 (13)
Pulmonary medicine	115 (8)
Other	87 (6)
Surgery	38 (3)
Hospital	
General hospital	522 (34)
Private hospital	197 (13)
University hospital	421 (28)
University affiliated general hospital	363 (24)
Other	18 (1)
Type of ICU	
Burn unit	4 (0.2)
Cardiac ICU	43 (3)
Coronary ICU	1 (0.1)
Mixed ICU	862 (57)
Medical ICU	312 (21)
Neurologic ICU	41 (3)
Surgical ICU	170 (11)
Trauma ICU	24 (2)
Transplant ICU	12 (1)
Other	52 (3)
ICU setting	
Closed ICU	841 (55)
Open ICU	183 (12)
Semiclosed	487 (32)
Other	10 (1)

(Continued)

TABLE 1. (Continued). Characteristics of 1,521 Responders in 47 Countries

Variables	n (%)
ICU no. of beds	
0–10	481 (32)
10–20	642 (42)
> 20	398 (26)
Patients admitted per yr	
301–500	442 (34)
501–1,000	431 (33)
> 1,000	353 (27)
Not available	78 (6)
Ventilated patients ^a (%)	
51–70	567 (37)
≤ 50	579 (38)
> 70	346 (23)
Not available	29 (2)
ICU length of stay (d)	
0–5	710 (47)
6–10	552 (36)
11–15	135 (9)
16–20	57 (4)
21–25	20 (1)
26–30	24 (2)
> 30	23 (2)
Nurse:patients ratio (d)	
1:1	164 (11)
1:2	976 (64)
1:3	270 (18)
1:4	111 (7)
Nurse:patients ratio (night)	
1:1	117 (8)
1:2	677 (44)
1:3	527 (35)
1:4	200 (13)

^aPercentage of patients receiving invasive mechanical ventilation over the total no. of ICU patients.

When delirium is identified, 74% of respondents would investigate potential causes, but significant heterogeneity was reported in the sequence of diagnostic methods used; the preferred combination was neurologic examination followed by a review of medications, laboratory tests, and infection screening. Respondents reported rarely using

neuroimaging, electroencephalography, and evoked potentials to investigate delirium. Just under half of respondents believe that delirium could affect mortality, ICU and hospital length of stay, ICU cost, family burden, or cognitive impairment.

Among nonpharmacologic interventions to promote sleep, respondents most commonly (28%) prefer optimizing ambient light, timing of drug administration, and noise. Alternatively, 58% of respondents reported prescribing drugs to promote sleep, with benzodiazepine alone (11%) being the preferred agent (data not shown).

When managing delirium, 58% of respondents reported that they do not use a protocol, and 65% reporting using haloperidol, most often (44%) as a single dose. Atypical antipsychotics are used by 53% respondents, either as a first approach (39%) or when haloperidol is not effective (42%). Respondents report consulting a specialist in 64% of delirium cases, usually (63%) in the most challenging cases.

Early Mobilization and Exercise

Just over one third of respondents reported that they routinely assess patients for ICU-acquired weakness (ICU-AW), most

TABLE 2. ABC: Assess, Prevent, and Manage Pain; Spontaneous Awakening Trials and Spontaneous Breathing Trials, Choice Of Analgesia and Sedation

Variables	Africa, n = 43 (%)	Asia, n = 441 (%)	Europe, n = 607 (%)	North America, n = 120 (%)	Oceania, n = 45 (%)	South America, n = 265 (%)	Total, n = 1,521 (%)
Assess, prevent, and manage pain							
Do you use protocol for pain treatment?							
No	31 (72)	114 (26)	256 (42)	45 (38)	29 (64)	110 (42)	585 (38)
Yes	12 (28)	327 (74)	351 (58)	75 (63)	16 (36)	155 (59)	936 (62)
Do you use any scale to evaluate pain in your unit?							
No	26 (60)	40 (9)	103 (17)	8 (7)	15 (33)	60 (23)	252 (16)
Yes	17 (40)	401 (91)	504 (83)	112 (93)	30 (67)	205 (77)	1269 (83)
Scale used to evaluate pain ^a							
Critical-Care Pain Observation Tool	0 (0)	46 (11)	52 (10)	70 (65)	6 (20)	41 (20)	215 (17)
Behavioral Pain Scale	1 (6)	140 (35)	129 (26)	14 (13)	1 (3)	53 (26)	338 (27)
Numerical Rating Scale	11 (65)	224 (56)	292 (58)	50 (45)	18 (60)	94 (46)	689 (54)
Visual Analogic Scale	5 (29)	272 (68)	284 (56)	21 (19)	12 (40)	86 (42)	680 (54)
Faces Pain Scale	7 (41)	80 (20)	99 (20)	19 (17)	9 (30)	39 (19)	253 (20)
Adult Nonverbal Pain Scale	2 (12)	7 (2)	52 (10)	19 (17)	3 (10)	17 (8)	100 (8)
Do you use preemptive analgesia before nursing procedure?							
No	32 (74)	154 (27)	92 (15)	13 (11)	2 (4)	69 (26)	362 (24)
Yes	6 (14)	370 (66)	315 (52)	29 (24)	11 (24)	124 (47)	855 (56)
At nurse's discretion	5 (12)	37 (7)	200 (33)	78 (65)	32 (71)	71 (27)	423 (28)
Drugs used for pain treatment ^a							
Fentanyl	7 (16)	386 (69)	402 (66)	118 (98)	44 (98)	242 (92)	1199 (79)
Hydromorphone	1 (2)	14 (2)	23 (4)	102 (85)	8 (18)	6 (2)	154 (10)
Morphine	21 (48)	310 (55)	523 (86)	84 (70)	41 (91)	203 (77)	1182 (78)
Methadone	4 (9)	4 (1)	42 (7)	20 (17)	14 (31)	59 (22)	143 (9)
Remifentanyl	4 (9)	112 (20)	376 (62)	1 (1)	8 (18)	68 (26)	569 (37)
Paracetamol	37 (86)	282 (50)	506 (83)	44 (37)	43 (96)	142 (54)	1054 (69)
Nonsteroidal anti-inflammatory drugs	37 (86)	291 (52)	327 (54)	54 (45)	24 (53)	86 (33)	819 (54)

(Continued)

TABLE 2. (Continued). ABC: Assess, Prevent, and Manage Pain; Spontaneous Awakening Trials and Spontaneous Breathing Trials, Choice Of Analgesia and Sedation

Variables	Africa, n = 43 (%)	Asia, n = 441 (%)	Europe, n = 607 (%)	North America, n = 120 (%)	Oceania, n = 45 (%)	South America, n = 265 (%)	Total, n = 1,521 (%)
SATs and SBTs							
Do you perform SAT?							
No	33 (77)	71 (16)	322 (53)	7 (6)	18 (40)	62 (23)	513 (34)
Yes	10 (24)	370 (84)	285 (47)	113 (94)	27 (60)	203 (77)	1008 (66)
How many times do you perform SAT?							
As many times as possible	1 (10)	51 (14)	52 (18)	10 (9)	5 (19)	27 (13)	146 (15)
Every nurse's shift	0 (0)	26 (7)	20 (7)	9 (8)	0 (0)	1 (1)	56 (6)
I perform a sedation minimization strategy	3 (30)	49 (13)	76 (27)	16 (14)	10 (37)	39 (19)	193 (19)
Once a day	6 (60)	241 (65)	134 (47)	73 (65)	9 (33)	133 (66)	596 (59)
Other	0 (0)	3 (1)	3 (1)	5 (4)	3 (11)	3 (2)	17 (2)
Do you perform SBT?							
No	27 (63)	71 (16)	304 (50)	4 (3)	32 (71)	58 (22)	496 (32)
Yes	16 (37)	370 (84)	303 (50)	116 (97)	13 (29)	207 (78)	1025 (67)
Choice of analgesia and sedation							
Scale used to evaluate sedation ^a							
Motor Activity Assessment Scale	2 (5)	(1)	2 (0.3)	1 (1)	1 (2)	1 (0.4)	11 (1)
None	33 (77)	25 (6)	100 (16)	0 (0)	1 (2)	3 (1)	162 (11)
Other	1 (2)	1 (0.2)	8 (1)	3 (3)	0 (0)	1 (0.4)	14 (1)
Richmond Agitation Sedation Scale	5 (12)	273 (62)	290 (48)	108 (90)	37 (82)	214 (81)	927 (61)
Ramsey	2 (5)	120 (27)	186 (31)	1 (1)	1 (2)	26 (10)	336 (22)
Symptom Assessment Scale	0 (0)	18 (4)	21 (3)	7 (6)	5 (11)	20 (8)	71 (5)
Do you follow protocol for sedation in your unit?							
No	36 (83)	87 (20)	312 (51)	20 (17)	22 (49)	59 (22)	536 (35)
Yes	7 (16)	354 (80)	295 (49)	100 (83)	23 (51)	206 (78)	985 (65)
If yes, have you adopted a minimal or no sedation strategy?							
No	3 (43)	34 (10)	31 (11)	9 (9)	5 (22)	20 (10)	102 (10)
Yes	4 (57)	320 (90)	264 (89)	91 (91)	18 (78)	186 (90)	883 (90)
Do you attempt to minimize the use of benzodiazepines in your patients?							
No	9 (21)	38 (9)	91 (15)	3 (3)	1 (2)	14 (5)	156 (10)
Yes	34 (79)	403 (91)	516 (85)	117 (98)	44 (98)	251 (95)	1365 (90)
In a restless and agitated patient who screened positive at delirium assessment, how do you approach the sedation strategy after clinical evaluation?							
Delirium first→pain→sedation	2 (5)	31 (7)	62 (10)	10 (8)	4 (9)	28 (11)	137 (9)
Delirium→sedation→pain	3 (7)	9 (2)	22 (4)	4 (3)	1 (2)	3 (1)	42 (3)
None	4 (9)	2 (0.5)	24 (0.4)	0 (0)	3 (7)	4 (2)	37 (2)
Other	0 (0)	2 (0.5)	2 (0.3)	4 (3)	3 (7)	0 (0)	11 (1)
Pain→delirium→sedation	13 (30)	177 (40)	247 (41)	62 (52)	14 (31)	159 (60)	672 (44)
Pain→sedation→delirium	13 (30)	171 (39)	170 (28)	31 (26)	18 (40)	52 (20)	455 (30)
Sedation→pain→delirium	8 (19)	49 (11)	80 (13)	9 (8)	2 (4)	19 (7)	167 (11)

SAT = spontaneous awakening trial, SBT = spontaneous breathing trial.

^aMultiple responses were possible.

TABLE 3. D: Delirium Monitoring and Management

Variables	Africa, n = 43 (%)	Asia, n = 441 (%)	Europe, n = 607 (%)	North America, n = 120 (%)	Oceania, n = 45 (%)	South America, n = 265 (%)	Total, n = 1,521 (%)
Do you routinely monitor delirium in your unit?							
Never	23 (54)	84 (19)	276 (45)	5 (4)	14 (31)	54 (20)	456 (30)
More than once a day	4 (9)	125 (28)	122 (20)	94 (78)	19 (42)	87 (33)	451 (30)
Once a day	16 (37)	232 (53)	209 (34)	21 (18)	12 (27)	124 (47)	614 (40)
Which tools do you routinely use to assess delirium?							
Confusion Assessment Method-ICU	6 (14)	310 (90)	259 (90)	131 (88)	28 (64)	205 (78)	536 (35)
Intensive Care Delirium Screening Checklist	0	15 (4)	23 (8)	13 (9)	0 (0)	1 (0.4)	102 (7)
None	36 (86)	18 (5)	5 (2)	5 (3)	16 (36)	56 (21)	883 (58)
Do you generally investigate the potential causes of delirium in your patients?							
No	7 (16)	101 (23)	199 (33)	18 (15)	10 (22)	54 (20)	389 (26)
Yes	36 (84)	340 (77)	408 (67)	102 (85)	35 (78)	211 (80)	1132 (74)
In your experience, how many patients have delirium in your ICU? (%)							
0–10	20 (47)	98 (22)	128 (21)	10 (8)	5 (11)	36 (14)	297 (20)
11–30	19 (44)	217 (49)	310 (51)	48 (40)	24 (53)	127 (48)	745 (50)
31–60	4 (9)	116 (26)	145 (24)	45 (38)	14 (31)	89 (34)	413 (27)
> 60	0 (0)	10 (2)	24 (4)	17 (14)	2 (4)	13 (5)	66 (4)
Do you prescribe earplugs to your patients?							
No	41 (95)	305 (69)	541 (89)	91 (76)	29 (64)	243 (92)	1250 (82)
Yes	2 (5)	136 (31)	66 (11)	29 (24)	16 (36)	22 (8)	271 (18)
Do you use nonpharmacologic interventions to promote sleep in your unit?							
Drug time optimization	4 (10)	30 (7)	47 (8)	4 (4)	2 (5)	34 (14)	121 (8)
Drug time optimization and noise reduction	1 (3)	5 (1)	30 (5)	6 (5)	0 (0)	6 (2)	48 (3)
Light optimization	6 (15)	118 (27)	72 (12)	10 (9)	1 (2)	42 (17)	249 (17)
Light optimization and drug time optimization	0 (0)	22 (5.1)	52 (9)	6 (5)	3 (7)	33 (13)	116 (8)
Light optimization, drug time optimization, and noise reduction	9 (23)	82 (19)	204 (35)	39 (34)	17 (42)	59 (23)	410 (28)
Light optimization and noise reduction	6 (15)	144 (33)	125 (21)	24 (21)	12 (30)	48 (19)	359 (24)
Noise reduction	13 (33)	31 (7)	62 (10)	26 (23)	6 (15)	30 (12)	168 (11)
Do you use protocols for the management of delirium?							
No	37 (86)	179 (41)	439 (71)	58 (48)	33 (73)	130 (49)	876 (58)
Yes	6 (14)	262 (59)	168 (28)	62 (52)	12 (27)	135 (51)	645 (42)

(Continued)

TABLE 3. (Continued). D: Delirium Monitoring and Management

Variables	Africa, n = 43 (%)	Asia, n = 441 (%)	Europe, n = 607 (%)	North America, n = 120 (%)	Oceania, n = 45 (%)	South America, n = 265 (%)	Total, n = 1,521 (%)
How do you use haloperidol to manage delirium in your unit?							
None	6 (14)	29 (7)	44 (7)	16 (13)	6 (13)	11 (4)	112 (7)
Prevention of delirium	2 (5)	2 (0.5)	13 (2)	0 (0)	0 (0)	3 (1)	20 (1)
Prevention of delirium + sedatives minimization	0 (0)	4 (1)	9 (2)	0 (0)	0 (0)	3 (1)	16 (1)
Sedatives minimization	2 (5)	20 (5)	40 (7)	6 (5)	0 (0)	8 (3)	76 (5)
Treatment of delirium episode	27 (63)	351 (80)	368 (61)	58 (48)	33 (73)	148 (56)	985 (65)
Treatment of delirium episode + prevention of delirium	1 (2)	10 (2)	36 (6)	3 (3)	2 (4)	10 (4)	62 (4)
Treatment delirium episode + prevention of delirium + sedatives minimization	2 (5)	9 (2)	31 (5)	6 (5)	0 (0)	16 (6)	64 (4)
Treatment of delirium episode + sedatives minimization	3 (7)	16 (4)	66 (11)	31 (26)	4 (9)	66 (25)	186 (12)
Do you use quetiapine or other atypical antipsychotic for delirium with agitation?							
No	36 (84)	248 (56)	342 (56)	14 (12)	15 (33)	63 (24)	718 (47)
Yes	7 (16)	193 (44)	265 (44)	106 (88)	30 (67)	202 (76)	803 (53)
In which situation do you require a specialist consultation (i.e., psychiatrist, neurologist, geriatrician)? ^a							
All cases	12 (28)	70 (16)	42 (7)	2 (3)	2 (4)	12 (5)	141 (9)
Most challenging cases of delirium	31 (72)	337 (76)	370 (61)	62 (52)	21 (47)	144 (54)	965 (63)
Alcohol withdrawal syndrome	14 (33)	203 (46)	73 (12)	9 (8)	4 (9)	51 (19)	354 (23)
Posttraumatic stress disorder	14 (33)	195 (44)	110 (18)	25 (21)	8 (18)	55 (21)	407 (27)
Never	2 (5)	12 (7)	125 (21)	39 (32)	17 (38)	68 (26)	263 (17)

^aMultiple responses were possible.

often using the Medical Research Council scale (49%) or an electrophysiologic evaluation (41%) (Table 4; and Appendix 5, Fig. 2, Supplemental Digital Content 2, <http://links.lww.com/CCM/C793>). Seventy three percent to 91% of the respondents from various continents report prescribing early mobilization, though they report rarely using a specific mobility scale (21%). Approaches reported consisted of combined passive range of motion, active physiotherapy, and ambulation (32%). Cycleergometry (14%) and neuromuscular electrical stimulation (6%) were infrequently reported. One third (31%) reported having a mobility team, consisting of a physical therapist (33%) or physical therapist and ICU nurses (17%), or physical therapist, respiratory therapist, and ICU nurses (12%) (Appendix 6, Supplemental Digital Content 2, <http://links.lww.com/CCM/>

C793). Sixty-five percent of the respondents who reported the use of early mobilization used specific scales to evaluate delirium.

Family

Of the respondents, 65% report that their unit is not open 24 hours per day for family visitation, 74% report that family member visits are allowed less than 5 hr/d (Table 5; and Appendix 5, Fig. 2, Supplemental Digital Content 2, <http://links.lww.com/CCM/C793>). Eighty-one percent of respondents report that they explain delirium to family members, with 13% reporting they use booklets. Family members are actively involved in 67% of the cases, but only 33% of the ICUs use dedicated staff to support families. When family members are actively involved, there is a higher prevalence

TABLE 4. E: Early Mobilization and Exercise

Variables	Africa, n = 43 (%)	Asia, n = 441 (%)	Europe, n = 607 (%)	North America, n = 120 (%)	Oceania, n = 45 (%)	South America, n = 265 (%)	Total, n = 1,521 (%)
Do you evaluate ICU-acquired muscle weakness in your unit?							
No	27 (63)	322 (73)	384 (63)	84 (70)	19 (42)	136 (51)	972 (64)
Yes	16 (37)	119 (27)	223 (37)	36 (30)	26 (58)	129 (49)	549 (36)
Total	43	441	607	120	45	265	1,521
Do you prescribe early mobilization to your patients?							
No	3 (7)	45 (10)	59 (10)	4 (3)	4 (9)	18 (7)	133 (9)
Only in nonventilated patients	1 (2)	29 (7)	65 (11)	9 (8)	8 (18)	24 (9)	136 (9)
Yes	39 (91)	367 (83)	483 (80)	107 (89)	33 (73)	223 (84)	1,252 (82)
Total	43	441	607	120	45	265	1,521
Do you use an ICU mobility scale for goal-directed early mobilization?							
No	31 (72)	326 (74)	466 (77)	58 (48)	36 (80)	179 (68)	1,096 (79)
Yes	9 (21)	70 (16)	82 (14)	58 (48)	5 (11)	68 (26)	292 (21)
Total	40	396	548	116	41	247	1,388
Do you have a dedicated mobility team in your ICU?							
No	38 (88)	346 (78)	402 (66)	83 (69)	31 (69)	152 (57)	1,052 (69)
Yes	5 (12)	95 (22)	205 (34)	37 (31)	14 (31)	113 (43)	469 (31)
Total	43	441	607	120	45	265	1,521

of interventions to reduce and treat delirium (**Appendix 7**, Supplemental Digital Content 2, <http://links.lww.com/CCM/C793>), and interestingly, this remained the same for early exercise and mobilization, independent of the nurse patient ratio (**Appendix 8**, Supplemental Digital Content 2, <http://links.lww.com/CCM/C793>).

DISCUSSION

This is the first worldwide survey to assess the knowledge and use of the ABCDEF bundle. Of 1,521 respondents in 47 countries, 57% reported implementing the bundle. The large majority (83%, 89%, and 70%) reported evaluating pain, sedation, and delirium in their ICU, though only 42% reporting using a validated delirium tool. Almost two thirds reported using SATs and SBTs. Most report prescribing early mobility, but few report having designated mobility teams. Though family members were reported to be actively involved in most ICUs, few had dedicated staff to help family or incorporate family in decision-making.

Though previous surveys and point prevalence studies have evaluated the use of the ABCDEF approach in the management of critically ill patients (6–8), our investigation is the first international survey to assess use of the full ACBDEF bundle

rather than focus on single components (27–33). A multi-center European Survey found that 80% of ICUs reported routinely monitoring pain, with 93% using a validated tool for pain assessment (27). The most frequently used pain score was the VAS (63%) followed by the NRS (57%). Alternatively, an Australian point prevalence study found that pain was assessed only in 46% of 569 patients in 41 ICUs (28). We found similar reported rates in Europe although we observed higher rates of pain monitoring in Oceania compared with the previous Australasian study, though this might be related to an overestimation of the actual assessment (28).

Our finding that 84% of respondents in Asia report using SATs and SBTs is consistent with data from a recent survey from India (29). Our findings regarding use of SATs and SBTs in Europe (47% and 50%) are also consistent with previously reported data (27).

Our responses from intensivists in Asia, Australia, and Europe are comparable with those in previous surveys assessing use of sedation scales (27–29). In an Australian point prevalence study, routine sedation scale use was recorded in 63% of invasively ventilated patients (28). In a European Survey, routine sedation monitoring was reported in 88% of the ICUs, with most reporting use of RASS (54%), Ramsay (27%), or SAS (6%) (27).

TABLE 5. F: Family Engagement and Empowerment

Variables	Africa, n = 43 (%)	Asia, n = 441 (%)	Europe, n = 607 (%)	North America, n = 120 (%)	Oceania, n = 45 (%)	South America, n = 265 (%)	Total, n = 1,521 (%)
Is your unit open 24 hr per day to family members visit?							
No	25 (58)	234 (53)	490 (81)	21 (18)	9 (20)	204 (77)	983 (65)
Yes	18 (42)	207 (47)	117 (19)	99 (82)	36 (80)	61 (23)	538 (35)
Total	43	441	607	120	45	265	1521
How many hours is your unit open to family members if not 24 hr per day? (hr)							
0–5	18 (72)	213 (91)	314 (64)	2 (10)	0 (0)	176 (87)	723 (74)
5–10	2 (8)	5 (2)	36 (7)	11 (52)	0 (0)	4 (2)	58 (6)
10–15	0 (0)	1 (0.4)	6 (1)	1 (5)	2 (22)	1 (0.5)	11 (1)
15–20	5 (20)	15 (6)	134 (27)	2 (10)	4 (44)	23 (11)	183 (19)
≥ 20	0 (0)	0 (0)	0 (0)	5 (24)	3 (33)	0 (0)	8 (1)
Do you generally explain to family members what delirium is?							
No	16 (37)	87 (20)	122 (20)	16 (13)	11 (24)	32 (12)	284 (19)
Yes	27 (63)	354 (80)	485 (80)	104 (87)	34 (76)	233 (88)	1,237 (81)
Total	43	441	607	120	45	265	1,521
Do you use booklets or training material to improve delirium knowledge among family members?							
No	41 (95)	392 (89)	549 (90)	74 (62)	37 (82)	236 (89)	1,329 (87)
Yes	2 (5)	49 (11)	58 (10)	46 (38)	8 (18)	29 (11)	192 (13)
Total	43	441	607	120	45	265	1,521
Do you involve family member in the delirium management?							
No	20 (47)	111 (25)	255 (42)	26 (22)	8 (18)	74 (28)	494 (33)
Yes	23 (53)	330 (75)	352 (58)	94 (78)	37 (82)	191 (72)	1,027 (67)
Total	43	441	607	120	45	265	1,521
Do you use dedicated staff for managing the relationship with family members?							
No	28 (65)	239 (54)	511 (84)	82 (68)	32 (71)	127 (48)	1,019 (67)
Yes	15 (35)	202 (46)	96 (16)	38 (32)	13 (29)	138 (52)	502 (33)
Total	43	441	607	120	45	265	1521

In a survey of Indian ICUs, 58% of respondents reported routinely monitoring sedation level, with the Ramsay Sedation Scale being most often used (56%) followed by RASS (19%). Nearly all respondents (95%) reported using midazolam for sedation,

followed by propofol (68%) and dexmedetomidine (60%); fentanyl was the most common analgesic agent used (47%) (29). In our study, 35% of respondents reported that they do not use sedation protocols, especially those in Africa, Europe, and Oceania.

A survey conducted by the Indian Society of Critical Care Medicine found that 35% of intensivists reported assessing for delirium, using validated scales in only 22% of the cases (most commonly the CAM-ICU) (29). Similarly, a multicenter European Survey found that only 56% of respondents reported screening patients for symptoms of delirium (27). In an Australian point prevalence study, routine assessment of delirium occurred in only 3% of patients (28). In our study, delirium evaluation was reported to be much higher than in previous reports in Asia (81%) and in Australia (69%) (27–29), which could reflect overestimation on the part of respondents but might also reflect a change in clinical practice. CAM-ICU was reported to be the most widely used delirium monitoring tool, though 58% of respondents reported that they do not use a tool, particularly those in Africa (86%), Oceania (36%), and South America (21%).

Though its efficacy remains in question (36), haloperidol is used to treat delirium and/or minimize the use of sedatives. Respondents to an European Survey reported that antipsychotics were the most frequently used agents for delirium treatment although it was not clear if this choice was related to treatment of psychotic symptoms or agitation (27).

Two-point prevalence studies across 116 German hospitals reported that only 8% of ventilated patients received out-of-bed mobility, and only 3% of patients in 38 Australian/New Zealand ICUs achieved sitting at the edge of the bed with none standing, transferring to chair or walking (30, 31). A recent point prevalence study across 42 US ICUs reported that 32% of adult patients with acute respiratory failure (and 26% of ventilated patients) received physical/occupational therapy (32). An Indian survey reported that higher mobilization levels (86% at the bed side, 70% to a wheel chair and 67% limited ambulation) (29). Our results suggest a discordance in intent and resources for mobilization. Although 91% of respondents report prescribing early mobilization, only 36% say that they evaluate for ICU-AW and 31% report having a dedicated mobility team. Additionally, there is significant variability in the composition of ICU mobility teams, with only a minority including a physical therapist, occupational therapist, nurses, and physicians.

Despite broad consensus that liberalization of visiting hours in the ICU improves the care and experience of patients and families (37, 38), a recent multicenter Brazilian survey reported that only 3% of the ICUs had liberal visitation policies (39). Among 289 French ICUs, only 24% were open for family visits 24 hours per day (40). Similarly, most of our respondents (65%) reported that their units are not open 24 hours per day, with most ICUs (74%) open from 5 hours per day or less. Despite evidence of the benefit of family engagement, it remains unclear how family involvement should be structured (41).

Our study has strengths and limitations. This is the first worldwide survey to explore the knowledge of the ABCDEF bundle and its use in clinical practice. The study provides detailed information about each component of ABCDEF, thus improving our understanding of the practices worldwide. The

precise response rate cannot be precisely determined due to the difficulties in conducting such a wide spread web-based survey, and there is a potential for selection bias due to the method of survey distribution. However, our survey respondents covered a broad range of age groups, clinical experience, and types of ICUs, including both teaching and nonteaching hospitals of various sizes. Consequently, our data reflect the broad spectrum of clinical practice across regions and estimates the range of current clinical practice. Additionally, our results suggest which elements of the bundle are the least implemented, thus providing targets for quality improvement initiatives, as well as those which need better infrastructural support for implementation.

This survey was limited to physicians; hence, we are unable to compare the responses with that of the other ICU staff, for example, nurses, physical therapists, who may have a different perspective and play an important role in the implementation of the ABCDEF bundle. An inherent limitation of “self-reporting” is that the reliability of individual responses cannot be ensured. There is also a possibility of having several participants from the same institution, which may result in multiple same or similar responses. Another limitation, which is not in our control, is the higher number of responses collected from single countries, potentially reflecting the care practices in those specific countries rather than the continent. Despite these limitations, this survey provides a fair idea about the knowledge and use of the ABCDEF bundle, which may help plan future research and improved strategies for implementation.

CONCLUSIONS

Just over decade ago, the majority of ICUs were closed to family members, practicing heavy sedation and patient immobilization. Our data reflect a dramatic yet incomplete cultural shift toward a patient- and family-centered ICU liberation strategy. There remains a compelling need for greater implementation of the ABCDEF bundle, particularly concerning the management of sedation, full appreciation assessment of delirium, and application of early mobility. An open ICU visitation policy is still rare, and there is a growing need to improve interaction with family members.

ACKNOWLEDGMENTS

We thank the following international critical care societies for their support:

- 1) The Australasian and New Zealand Intensive Care Society.
- 2) The European Society of Intensive Care Medicine.
- 3) The Italian Society of Anesthesia, Analgesia, Reanimation and Intensive Care.
- 4) The Indian Society of Critical Care Medicine.
- 5) The Japanese Society of Critical Care Medicine.

REFERENCES

1. Brummel NE, Bell SP, Girard TD et al: Frailty and subsequent disability and mortality among patients with critical illness. *Am J Respir Crit Care Med.* 2017; 196:64–72

2. Pandharipande PP, Girard TD, Jackson JC, et al; BRAIN-ICU Study Investigators: Long-term cognitive impairment after critical illness. *N Engl J Med* 2013; 369:1306–1316
3. Wolters AE, Peelen LM, Veldhuijzen DS, et al: Long-term self-reported cognitive problems after delirium in the intensive care unit and the effect of systemic inflammation. *J Am Geriatr Soc* 2017; 65:786–791
4. Jackson JC, Pandharipande PP, Girard TD, et al; Bringing to light the Risk Factors And Incidence of Neuropsychological dysfunction in ICU survivors (BRAIN-ICU) study investigators: Depression, post-traumatic stress disorder, and functional disability in survivors of critical illness in the BRAIN-ICU study: A longitudinal cohort study. *Lancet Respir Med* 2014; 2:369–379
5. Latronico N, Bolton CF: Critical illness polyneuropathy and myopathy: a major cause of muscle weakness and paralysis. *Lancet Neurol* 2011; 10:931–941
6. Balas MC, Burke WJ, Gannon D, et al: Implementing the awakening and breathing coordination, delirium monitoring/management, and early exercise/mobility bundle into everyday care: Opportunities, challenges, and lessons learned for implementing the ICU Pain, Agitation, and Delirium Guidelines. *Crit Care Med* 2013; 41(9 Suppl 1):S116–S127
7. Morandi A, Brummel NE, Ely EW: Sedation, delirium and mechanical ventilation: The 'ABCDE' approach. *Curr Opin Crit Care* 2011; 17:43–49
8. Vasilevskis EE, Pandharipande PP, Girard TD, et al: A screening, prevention, and restoration model for saving the injured brain in intensive care unit survivors. *Crit Care Med* 2010; 38(10 Suppl):S683–S691
9. Barnes-Daly MA, Phillips G, Ely EW: Improving hospital survival and reducing brain dysfunction at seven California community hospitals: Implementing PAD guidelines via the ABCDEF bundle in 6,064 patients. *Crit Care Med* 2017; 45:171–178
10. Girard TD, Kress JP, Fuchs BD, et al: Efficacy and safety of a paired sedation and ventilator weaning protocol for mechanically ventilated patients in intensive care (Awakening and Breathing Controlled trial): A randomised controlled trial. *Lancet* 2008; 371:126–134
11. Kress JP, Pohlman AS, O'Connor MF, et al: Daily interruption of sedative infusions in critically ill patients undergoing mechanical ventilation. *N Engl J Med* 2000; 342:1471–1477
12. Kress JP, Gehlbach B, Lacy M, et al: The long-term psychological effects of daily sedative interruption on critically ill patients. *Am J Respir Crit Care Med* 2003; 168:1457–1461
13. Mehta S, Burry L, Martinez-Motta JC, et al; Canadian Critical Care Trials Group: A randomized trial of daily awakening in critically ill patients managed with a sedation protocol: A pilot trial. *Crit Care Med* 2008; 36:2092–2099
14. Pandharipande PP, Sanders RD, Girard TD, et al; MENDS investigators: Effect of dexmedetomidine versus lorazepam on outcome in patients with sepsis: An a priori-designed analysis of the MENDS randomized controlled trial. *Crit Care* 2010; 14:R38
15. Pandharipande PP, Pun BT, Herr DL, et al: Effect of sedation with dexmedetomidine vs lorazepam on acute brain dysfunction in mechanically ventilated patients: The MENDS randomized controlled trial. *JAMA* 2007; 298:2644–2653
16. Jakob SM, Ruokonen E, Grounds RM, et al; Dexmedetomidine for Long-Term Sedation Investigators: Dexmedetomidine vs midazolam or propofol for sedation during prolonged mechanical ventilation: Two randomized controlled trials. *JAMA* 2012; 307:1151–1160
17. Riker RR, Shehabi Y, Bokesch PM, et al; SEDCOM (Safety and Efficacy of Dexmedetomidine Compared With Midazolam) Study Group: Dexmedetomidine vs midazolam for sedation of critically ill patients: A randomized trial. *JAMA* 2009; 301:489–499
18. Fraser GL, Devlin JW, Worby CP, et al: Benzodiazepine versus non-benzodiazepine-based sedation for mechanically ventilated, critically ill adults: A systematic review and meta-analysis of randomized trials. *Crit Care Med* 2013; 41(9 Suppl 1):S30–S38
19. Zhang Z, Pan L, Ni H: Impact of delirium on clinical outcome in critically ill patients: A meta-analysis. *Gen Hosp Psychiatry* 2013; 35:105–111
20. Salluh JJ, Wang H, Schneider EB, et al: Outcome of delirium in critically ill patients: Systematic review and meta-analysis. *BMJ* 2015; 350:h2538
21. Patel SB, Poston JT, Pohlman A, et al: Rapidly reversible, sedation-related delirium versus persistent delirium in the intensive care unit. *Am J Respir Crit Care Med* 2014; 189:658–665
22. Schweickert WD, Pohlman MC, Pohlman AS, et al: Early physical and occupational therapy in mechanically ventilated, critically ill patients: A randomised controlled trial. *Lancet* 2009; 373:1874–1882
23. Li Z, Peng X, Zhu B, et al: Active mobilization for mechanically ventilated patients: A systematic review. *Arch Phys Med Rehabil* 2013; 94:551–561
24. Chen YH, Lin HL, Hsiao HF, et al: Effects of exercise training on pulmonary mechanics and functional status in patients with prolonged mechanical ventilation. *Respir Care* 2012; 57:727–734
25. Morris PE, Goad A, Thompson C, et al: Early intensive care unit mobility therapy in the treatment of acute respiratory failure. *Crit Care Med* 2008; 36:2238–2243
26. Schaller SJ, Anstey M, Blobner M, et al; International Early SOMS-guided Mobilization Research Initiative: Early, goal-directed mobilisation in the surgical intensive care unit: A randomised controlled trial. *Lancet* 2016; 388:1377–1388
27. Luetz A, Balzer F, Radtke FM, et al: Delirium, sedation and analgesia in the intensive care unit: A multinational, two-part survey among intensivists. *PLoS One* 2014; 9:e110935
28. Elliott D, Aitken LM, Bucknall TK, et al; Australian and New Zealand Intensive Care Society Clinical Trials Group; George Institute for Global Health: Patient comfort in the intensive care unit: A multicentre, binational point prevalence study of analgesia, sedation and delirium management. *Crit Care Resusc* 2013; 15:213–219
29. Chawla R, Myatra SN, Ramakrishnan N, et al: Current practices of mobilization, analgesia, relaxants and sedation in Indian ICUs: A survey conducted by the Indian Society of Critical Care Medicine. *Indian J Crit Care Med* 2014; 18:575–584
30. Nydahl P, Ruhl AP, Bartoszek G, et al: Early mobilization of mechanically ventilated patients: A 1-day point-prevalence study in Germany. *Crit Care Med* 2014; 42:1178–1186
31. Berney SC, Harrold M, Webb SA, et al: Intensive care unit mobility practices in Australia and New Zealand: A point prevalence study. *Crit Care Resusc* 2013; 15:260–265
32. Jolley SE, Moss M, Needham DM, et al; Acute Respiratory Distress Syndrome Network Investigators: Point prevalence study of mobilization practices for acute respiratory failure patients in the United States. *Crit Care Med* 2017; 45:205–215
33. Pinto F, Biancofiore G: The ABCDE bundle: A survey of nurses knowledge and attitudes in the intensive care units of a national teaching hospital in Italy. *Dimens Crit Care Nurs* 2016; 35:309–314
34. Shinotsuka CR, Salluh JJ: Perceptions and practices regarding delirium, sedation and analgesia in critically ill patients: A narrative review. *Rev Bras Ter Intensiva* 2013; 25:155–161
35. Eysenbach G: Improving the quality of Web surveys: The Checklist for Reporting Results of Internet E-Surveys (CHERRIES). *J Med Internet Res* 2004; 6:e34
36. Neufeld KJ, Yue J, Robinson TN, et al: Antipsychotic medication for prevention and treatment of delirium in hospitalized adults: A systematic review and meta-analysis. *J Am Geriatr Soc* 2016; 64:705–714
37. Giannini A, Garrouste-Orgeas M, Latour JM: What's new in ICU visiting policies: Can we continue to keep the doors closed? *Intensive Care Med* 2014; 40:730–733
38. Azoulay E, Pochard F, Chevret S, et al; French Famirea Group: Family participation in care to the critically ill: Opinions of families and staff. *Intensive Care Med* 2003; 29:1498–1504
39. Ramos FJ, Fumis RR, de Azevedo LC, et al: Intensive care unit visitation policies in Brazil: A multicenter survey. *Rev Bras Ter Intensiva* 2014; 26:339–346
40. Garrouste-Orgeas M, Vinatier I, Tabah A, et al: Reappraisal of visiting policies and procedures of patient's family information in 188 French ICUs: A report of the Outcomerea Research Group. *Ann Intensive Care* 2016; 6:82
41. Liput SA, Kane-Gill SL, Seybert AL, et al: A review of the perceptions of healthcare providers and family members toward family involvement in active adult patient care in the ICU. *Crit Care Med* 2016; 44:1191–1197